

NOTIFICATION TO ALL SURGICAL PATIENTS

PLEASE BRING A LIST OF YOUR MEDICATIONS AND ALLERGIES AT PRE-OP.

Please discuss with the staff which medications need to be continued or held prior to your surgery.

There are certain medications (**over the counter and prescribed**) that may cause **BLEEDING**. These medications need to be stopped 2 weeks prior to your surgical procedure. Do not resume these medications until cleared by the clinical staff. **TYLENOL** does not cause bleeding and may be taken for pain before your surgery.

AVOID the following medications:

Advil
Alka-Seltzer
Aleve
Anacin
Anaprox
A.P.C.
Apectol Tabs
Arthritis Strength Bufferin
Ascodeen-30
Ascription
ASPIRIN
Aspirin Suppositories
BC Powder
Buff-a-Comp
Buffadyne
Bufferin
Butelbital
Cama-Inlay Tab
Cataflam
Clinoril
Congespirin
Cope Coricidin
Damason-P
Darvon Compound
Darvon w/Aspirin
Darvon-N
Daypro
Dristan
Duragesic

Easprin
Ecotrin
Emprazol
Empirin
Empirin w/Codeine
Equagesic
Excedrin
Excedrin X-Strength
Extra Strength Doans
Feldene
Fiorinal
4-Way Cold tabs
Ger-o-Form
Goodies

Haltran Tabs
Ibuprofen
Indocin
Lortab ASA
Medigesic Plug
Measurin
Midol
Mobigesic
Monocet w/Codeine
Motrin
Naprosyn
Norgesic
Nuprin
Orudis
Oruvail

Os-Cal-Gesic
Pabrinin Buffered Tabs
Panalfesic
Peda-Profen
Percodan
Persantin
Plavix
Ponstel
Relafen
Rhinex
Robaxsial
Rufen
Salsalate Disalcid
Sine-Aid
Sine-Off
SK-65 Comp. Caps
Soma Compound
Stendin
Supac
Stero-Darvon w/Aspirin
Synalgos D.C.
Tabloid Brand A.P.C.
Tolectin
Toradol
Triaminicin
Trilisnte
Vanquish
Voltaren

Warfarin/Coumadin

STOP all ARTHRITIC MEDICATIONS, VITAMIN E, and HERBAL REMEDIES

CONTINUE the following medications: Heart/Blood Pressure Medications

If you are **Diabetic discuss your medications and glucose levels at pre-op with the nurse.*