

NOTIFICATION TO ALL SURGICAL PATIENTS

PLEASE BRING A LIST OF YOUR MEDICATIONS AND ALLERGIES AT PRE-OP.

Please discuss with the staff which medications need to be continued or held prior to your surgery.

There are certain medications (**over the counter and prescribed**) that may cause **BLEEDING**. These medications need to be stopped 2 weeks prior to your surgical procedure. Do not resume these medications until cleared by the clinical staff. **TYLENOL** does not cause bleeding and may be taken for pain before your surgery.

AVOID the following medications:

Advil	Easprin	Os-Cal-Gesic
Alka-Seltzer	Ecotrin	Pabrinin Buffered Tabs
Aleve	Empirazil	Panalfesic
Anacin	Empirin	Peda-Profen
Anaprox	Empirin w/Codeine	Percodan
A.P.C.	Equagesic	Persantin
Apectol Tabs	Excedrin	Plavix
Arthritis Strength Bufferin	Excedrin X-Strength	Ponstel
Ascodeen-30	Extra Strength Doans	Relafen
Ascription	Feldene	Rhinex
<u>ASPIRIN</u>	Fiorinal	Robaxsial
Aspirin Suppositories	4-Way Cold tabs	Rufen
BC Powder	Ger-o-Form	Salsalate Disalcid
Buff-a-Comp	Goodies	Sine-Aid
Buffadyne	Haltran Tabs	Sine-Off
Bufferin	Ibuprofen	SK-65 Comp. Caps
Butelbital	Indocin	Soma Compound
Cama-Inlay Tab	Lortab ASA	Stendin
Cataflam	Medigesic Plug	Supac
Clinoril	Measurin	Stero-Darvon w/Aspirin
Congespirin	Midol	Synalgos D.C.
Cope Coricidin	Mobigesic	Tabloid Brand A.P.C.
Damason-P	Monocet w/Codeine	Tolectin
Darvon Compound	Motrin	Toradol
Darvon w/Aspirin	Naprosyn	Triaminicin
Darvon-N	Norgesic	Trilisnte
Daypro	Nuprin	Vanquish
Dristan	Orudis	Voltaren
Duragesic	Oruvail	Warfarin/Coumadin

STOP all ARTHRITIC MEDICATIONS, VITAMIN E, and HERBAL REMEDIES

CONTINUE the following medications: **Heart/Blood Pressure Medications**

If you are **Diabetic discuss your medications and glucose levels at pre-op with the nurse.*