

# lachele

AESTHETICS & WELLNESS

## Filler Reversal - Hylenex Post Instructions

The treated area **may look swollen and non-dissolved** upon leaving our facility. Allow up to 2 weeks for full dissolution.

If intense swelling please immediately take Benadryl. If more severe swelling please call the office to see if you are a candidate for steroids.

Many **reversals can take 2-3 treatment** in a regular dermal filler. With Vycross technology (*Voluma, Vollure, Volbella*) or a granuloma case, **patients could require 3- 10 reversal sessions depending on the severity**. These reversals should be spaced 2-3 weeks apart depending on bruising.

As discussed during your consent, **dissolving can cause swelling and bruising** which can take 2 weeks or longer to resolve. **Apply ice in the first few days** and **then heat to a bruise post 5 days** if still present.

As discussed during your consent, the dissolving process will result in a loss of volume. This may result in the treatment area appearing more hollow or dark. **Re-filling cannot take place for at least 2 weeks post dissolving.**

As discussed during your consent, **future filler may be needed for restoration** of volume after the Hyaluronidase has settled. We cannot fully determine how the area will look post treatment or how much filler you may require to have a satisfactory result.

It is not guaranteed that adjacent areas to injection site will not be affected by the reversal agent.

The use of Hyaluronidase can result in anaphylaxis which is a severe allergic reaction that is life threatening. If you are experiencing any unusual symptoms such as shortness of breath or trouble breathing, please seek immediate medical attention.

If you have any questions or concerns, please call our offices at (855)522-4353 and follow the prompt for your desired location. During off hours, call Dr. Espinoza's cell at (908) 303-5694. You can also email us at [team@lachele.com](mailto:team@lachele.com)